



## Product Datasheet

<b>Product Name</b>	Regenerating Protein 1 beta Human Recombinant
<b>Cata No</b>	CB501012
<b>Source</b>	<i>Escherichia Coli.</i>
<b>Synonyms</b>	Lithostathine-1-beta, Regenerating protein I beta, REG1B, REGL.

### Description

Reg protein was shown to be stimulated during the regeneration of pancreatic islets. Since then, many Reg-related proteins have been identified in humans and other animals. In human, the four REG family genes, i.e., REG 1 alpha, REG 1 beta, REG-related sequence (RS) and HIP/PAP, have so far been isolated. These Reg-related proteins are classified into four subfamilies according to their amino-acid sequences, but they share a similar structure and physiological function. Reg protein is a growth factor for pancreatic beta cells and also suggests that the administration of Reg protein could be used as another therapeutic approach for diabetes mellitus. Human REG cDNA which encodes a 166-amino acid protein with a 22-amino acid signal peptide. The amino acid sequence of human REG protein has 68% homology to that of rat Reg protein. Reg I was found to be expressed mainly in pancreatic beta and acinoductular cells as well as gastric fundic enterochromaffin-like (ECL) cells. Reg I production in ECL cells is stimulated by gastrin, as well as by the proinflammatory cytokine, cytokine-induced neutrophil chemoattractant (CINC)-2Beta. In patients with chronic hypergastrinemia, Reg production is stimulated, with the increased proliferation of gastric mucosal cells. Patients with *Helicobacter pylori* infection also showed increased Reg production in the gastric mucosa, partly via increased plasma gastrin concentration and partly via increased proinflammatory cytokine production. The serum

concentration of the reg-protein was significantly higher in patients with various pancreatic diseases than in normal controls, and was also significantly higher in patients with acute pancreatitis or chronic relapsing pancreatitis than in patients with chronic pancreatitis. Furthermore, the serum PSP/reg-protein concentration was also significantly increased in liver cirrhosis, choledocholithiasis, and various cancers of the digestive system.

The Recombinant Human REG 1 beta manufactured with N-terminal fusion of His Tag. The Human REG 1 beta His-Tagged Fusion Protein, produced in *E. coli*, is 17.8 kDa protein containing 144 amino acid residues of the Human REG 1 beta and 12 additional amino acid residues – His Tag (underlined).

MKHHHHHHAS HMQESQTELP NPRISCPEGT  
NAYRSYCYFF NEDPETWVDA DLYCQNMNSG  
NLVSVLTQAE GAFVASLIKE SSTDDSNVWI  
GLHDPKKNRR WHWSSGSLVS YKSWDTGSPS  
SANAGYCASL TSCSGFKKWK DESCEKKFSF  
VCKFKN

### Purity

Greater than 95% as determined by SDS-PAGE.

### Formulation

Sterile filtered and lyophilized from 0.5 mg/ml in 20mM Tris, pH 8.0.

### Reconstitution

Add 0.2 ml of deionized H<sub>2</sub>O and let the lyophilized pellet dissolve completely.

**\* For Non-Clinical Research Use Only \***



# California Bioscience

83103 Avenue 48, Ste.1B #204

Coachella, CA 92236 USA

Phone : +1.6268339877

Email : [info@cali-bio.com](mailto:info@cali-bio.com)

## Stability

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to **avoid repeated freezing/thawing cycles**. Reconstituted protein can be stored at 4°C for a limited period of time; it does

not show any change after 12 weeks.

## Product Data sheet

The lyophilized protein remains stable until the expiry date when stored at -20°C.

## Applications

Western blotting, ELISA

**\* For Non-Clinical Research Use Only \***